

# Open Incisional Hernia Repair



An Open Incisional Hernia repair is an operation to mend a defect in the abdominal wall that has occurred after a surgical incision. The defect appears where the body tissues have failed to heal completely. With time a visible and often painful bulge then forms. The repair is achieved by the insertion of nylon mesh below, within or on top of the abdominal wall. The operation is a procedure performed under general anaesthetic.

An incision is made in the abdominal wall over the hernia, the edges of the hernia defect are cleared and the hernia contents (often bowel or fat) are pushed back into the abdominal cavity. The layers of the abdominal wall may be separated to allow the insertion of a mesh (usually of nylon) that will comfortably overlap the defect. The mesh is then fixed in position using tacks or stitches and the layers of the abdominal wall repaired in a similar fashion. The operation is sometimes performed as a day case but more usually with an overnight stay. A plastic tube drain may be placed within the repair to remove fluid. It will be taken out a few days later.

## What happens before the operation?

At a pre-assessment visit you will be seen by a doctor or nurse. You will have specific tests to ensure your fitness for surgery, which will depend upon your general medical health. The purpose of this visit is to spot any abnormalities well in advance of surgery and to make sure that there are no delays prior to timely, successful completion of your operation.

## What happens just before the operation?

You will be admitted to a surgical unit. A senior nurse will complete the admission process detailing your current condition, ongoing medical problems and medications as well as personal details such as next of kin. You will be permitted to eat up until six hours before surgery and drink clear fluids at the discretion of the doctors, usually up to two hours before the operation. An Anaesthetist will visit you prior to your surgery to discuss the nature of your anaesthetic and methods of pain control both during and after the operation. Your surgeon will visit you and discuss the operation again, asking you to sign a consent form once you are happy that all your questions have been answered and the risks and benefits have been explained to you.

## What are the complications of the operation?

Although there are infrequent risks attached to this operation the nature of the surgery means that these may be significant.

**Infection – Wounds and Mesh**  
The nature of the surgery means that infections are uncommon. They are usually no more complicated than a skin infection (usually around the port sites) which is easily treated with antibiotics. Mesh infection in all types of hernia repair is very uncommon, should it occur, reoperation and removal of the mesh may be required.

### Bleeding

Hernia surgery involves the division of many minor blood vessels. Although meticulous care is taken to prevent bleeding it is a recognised but small risk of such operations. It may require a return to the operating theatre but more usually a small haematoma (or bruise) develops which resolves with time.

### Recurrence

All types of hernia repair are measured by their recurrence rate. This is usually described as a percentage of all repairs that recur over a five year period after operation. Modern open hernia repair techniques even with mesh have a quoted recurrence rate of around 25%.

### Injury to other Structures

Since this operation is performed for patients who have previously had abdominal surgery, adhesions and as a result involvement of other structures is common. Damage to these structures is uncommon but may be significant or even life threatening should it occur.

### What will happen after the operation?

- The wound on your abdomen will be closed with absorbable stitches and/or paper strips and covered with a dressing. Dressings should be left in place for 5 days then removed. Usually they do not need to be replaced.
- You will be able to start drinking and eating immediately after surgery.
- You will be given a combination of pain killers by mouth on waking from surgery. Stronger intravenous painkillers are always available if required.
- Some discomfort after surgery is normal but pain is usually minimal and treated easily. Most patients take regular pain relief for only about a week.
- An abdominal support binder should be worn after surgery usually for two weeks.

### What happens next?

You may stay in hospital overnight or be able to go home the same day as long as you have no significant pain or nausea. A capable adult must remain with you for 24 hours after the operation.

### What happens after hospital?

How soon you go home after the operation depends on how quickly you recover from surgery and who is at home to help you. Before discharge you will be advised how to care for your wounds. There are no limitations on showering. Bathing should be avoided for a week. Stitches are absorbable and may be ignored. Sticky paper strips may also be used; these will fall off after a few days. Most people are largely back to normal at two weeks after such surgery but it may take up to six weeks before you can say that you are fully recovered from the operation. As you leave hospital your General Practitioner will be informed of your treatment and the plans for your follow up. You may be given or sent an appointment to be seen in the outpatient department by your surgeon between two and six weeks after discharge.

### How will I feel when I go home?

Initially it is not unusual to feel a little tired following your operation. Discomfort should be minimal and controlled with simple pain killers used in combination. You will be prescribed these on discharge. If you feel that pain is increasing around your wound then you should contact your surgeon. It is important to rest initially when getting home, but it is equally sensible to slowly increase your level of activity. Listen to your body and stop when you begin to tire. You are advised not to do any heavy lifting for six weeks after the operation. Driving should be avoided for at least a week. The main area of concern is your ability to perform an emergency stop. Check with your car insurance company regarding the cover you have following an operation.